



## Complete Summary

---

### GUIDELINE TITLE

Initial evaluation of the patient with lung cancer: symptoms, signs, laboratory tests, and paraneoplastic syndromes.

### BIBLIOGRAPHIC SOURCE(S)

Beckles MA, Spiro SG, Colice GL, Rudd RM. Initial evaluation of the patient with lung cancer: symptoms, signs, laboratory tests, and paraneoplastic syndromes. Chest 2003 Jan;123(1 Suppl):97S-104S. [54 references] [PubMed](#)

## COMPLETE SUMMARY CONTENT

SCOPE  
METHODOLOGY - including Rating Scheme and Cost Analysis  
RECOMMENDATIONS  
EVIDENCE SUPPORTING THE RECOMMENDATIONS  
BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS  
IMPLEMENTATION OF THE GUIDELINE  
INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT  
CATEGORIES  
IDENTIFYING INFORMATION AND AVAILABILITY

## SCOPE

### DISEASE/CONDITION(S)

Lung cancer

### GUIDELINE CATEGORY

Diagnosis  
Evaluation  
Risk Assessment  
Screening

### CLINICAL SPECIALTY

Oncology  
Pulmonary Medicine

### INTENDED USERS

Physicians

## GUIDELINE OBJECTIVE(S)

To provide clinically relevant evidence-based guidelines for the initial evaluation of patients with either known or suspected lung cancer

## TARGET POPULATION

Patients known or suspected to have lung cancer

## INTERVENTIONS AND PRACTICES CONSIDERED

### Evaluation

1. Chest x-ray
2. Evaluation of symptoms and signs related to the primary tumor
  - Cough
  - Dyspnea
  - Hemoptysis
  - Chest discomfort
3. Evaluation of signs and symptoms of intrathoracic spread
  - Recurrent laryngeal nerve palsy
  - Phrenic nerve paralysis
  - Pancoast tumor
  - Horner's syndrome
  - Chest wall pain
  - Pleuritic chest pain
  - Superior vena cava obstruction
  - Heart and pericardium metastasis
  - Esophageal metastasis
4. Evaluation of signs and symptoms of extrathoracic spread
  - Bone pain
  - Metastases to liver, adrenal glands, and intra-abdominal lymph nodes
  - Brain and spinal cord metastases
  - Lymph node and skin metastases
5. Standardized evaluation for systemic metastases
  - Symptoms (weight loss, focal skeletal pain, neurological symptoms)
  - Signs (lymphadenopathy, hoarseness, superior vena cava syndrome, bone tenderness, hepatomegaly, focal neurological signs, papilledema, soft-tissue mass)
  - Laboratory tests (hematocrit, alkaline phosphatase, gamma-glutamyltransferase, serum glutamicoxaloacetic acid)
6. Evaluation for paraneoplastic syndromes
  - Endocrine paraneoplastic syndromes (e.g., hypercalcemia, syndrome of inappropriate antidiuretic syndrome [SIADH], Cushing syndrome)
  - Digital clubbing and hypertrophic osteoarthropathy
  - Neurologic syndromes

## MAJOR OUTCOMES CONSIDERED

- Frequencies of local and systemic symptoms in patients with lung cancer
- Frequencies of paraneoplastic syndromes in patients with lung cancer
- 5-year survival

## METHODOLOGY

### METHODS USED TO COLLECT/SELECT EVIDENCE

Hand-searches of Published Literature (Primary Sources)  
Searches of Electronic Databases

### DESCRIPTION OF METHODS USED TO COLLECT/SELECT THE EVIDENCE

As a first step in identifying the evidence for each topic, the guideline developers sought existing evidence syntheses including guidelines, systematic reviews, and meta-analyses. They searched computerized bibliographic databases including MEDLINE, Cancerlit, CINAHL and HealthStar, the Cochrane Collaboration Database of Abstracts of Reviews of Effectiveness, the National Guideline Clearinghouse, and the National Cancer Institute Physician Data Query database. Computerized searches through July 2001 used the MeSH terms lung neoplasms (exploded) and bronchial neoplasms or text searches for lung cancer combined with review articles, practice guidelines, guidelines, and meta-analyses. They also searched and included studies from the reference lists of review articles, and queried experts in the field. An international search was conducted of Web sites of provider organizations that were likely to have developed guidelines. Abstracts of candidate English language articles were reviewed by two physicians (one with methodological expertise and one with content area expertise) and a subset was selected for review in full text. Full-text articles were reviewed again by two physicians to determine whether they were original publications of a synthesis and were pertinent to at least one of the topics of the guideline. Articles described as practice guidelines, systematic reviews, or meta-analyses were included, as were review articles that included a "Methods" section. Included articles were classified according to topic.

### NUMBER OF SOURCE DOCUMENTS

Not stated

### METHODS USED TO ASSESS THE QUALITY AND STRENGTH OF THE EVIDENCE

Expert Consensus  
Weighting According to a Rating Scheme (Scheme Given)

### RATING SCHEME FOR THE STRENGTH OF THE EVIDENCE

The United States Preventive Services Task Force (USPSTF) scheme offers general guidelines to assign one of the following grades of evidence: good, fair, or poor. In general, good evidence included prospective, controlled, randomized clinical trials, and poor evidence included case series and clinical experience. Trials with fair quality of evidence, for instance, historically controlled trials or retrospective analyses, were somewhere in between. In addition to the strength of the study design, however, study quality also was considered. The USPSTF approach considers well-recognized criteria in rating the quality of individual studies for a variety of different types of study design (e.g., diagnostic accuracy studies and

case-control studies). The thresholds for distinguishing good vs fair and fair vs poor evidence are not explicit but are left to the judgment of panelists, reviewers, and members of the executive committee.

## Assessment of the Scope and Quality of Clinical Practice Guidelines

Clinical practice guidelines identified from the systematic search were evaluated by at least four reviewers using the Appraisal of Guidelines for Research and Evaluation (AGREE) instrument.

## METHODS USED TO ANALYZE THE EVIDENCE

Review of Published Meta-Analyses  
Systematic Review

## DESCRIPTION OF THE METHODS USED TO ANALYZE THE EVIDENCE

Not stated

## METHODS USED TO FORMULATE THE RECOMMENDATIONS

Informal Consensus

## DESCRIPTION OF METHODS USED TO FORMULATE THE RECOMMENDATIONS

Each writing committee received a comprehensive list of existing systematic reviews and meta-analyses as well as guidelines published by other groups. In addition, for five key topics (prevention, screening, diagnosis, and staging [invasive and noninvasive], new systematic reviews were undertaken [see "Description of Methods Used to Collect the Evidence" and "Description of Methods Used to Analyze the Evidence" fields]). For all other topics, writing committees were responsible for identifying and interpreting studies that were not otherwise covered in existing syntheses or guidelines.

The guidelines developed by the writing committee were distributed to the entire expert panel, and comments were solicited in advance of a meeting. During the meeting, proposed recommendations were reviewed, discussed, and voted on by the entire panel. Approval required consensus, which was defined as an overwhelming majority approval. Differences of opinion were accommodated by revising the proposed recommendation, the rationale, or the grade until consensus could be reached. The evidence supporting each recommendation was summarized, and recommendations were graded as described. The assessments of level of evidence, net benefit, and grade of recommendation were reviewed by the executive committee.

## Values

The panel considered data on functional status, quality and length of life, tolerability of treatment, and relief of symptoms in formulating guideline recommendations. Cost was not explicitly considered in the guideline development

process. Data on these outcomes were informally weighted, without the use of explicit decision analysis or other modeling. The values placed on types of outcomes varied with clinical scenarios. For example, in some situations they considered life expectancy, such as the effects of early detection. In other situations they weighed quality of life more heavily, such as in palliative care and in interpreting small increases in life expectancy with chemotherapy for stage IV disease.

## RATING SCHEME FOR THE STRENGTH OF THE RECOMMENDATIONS

The guideline developer's grading scheme is a modification of the United States Preventive Services Task Force (USPSTF) grades to allow recommendations for a service when (1) evidence is poor, (2) the assessment of the net benefit is moderate to high, and (3) there is consensus among the expert panel to recommend it. This change was necessary because, unlike preventive services (i.e., the routine offering of tests or treatments to well people) in which the burden of proof is high, clinical decisions about the treatment of patients with lung cancer often must be based on an interpretation of the available evidence, even if it is of poor quality. This adaptation distinguished between interventions with poor evidence for which there is consensus (grade C) and interventions with poor evidence for which there is not consensus (grade I).

### Grades of Recommendations and Estimates of Net Benefit

The grade of the strength of recommendations is based on both the quality of the evidence and the net benefit of the service (i.e., test, procedure, etc).

**Grade A** The panel strongly recommends that clinicians routinely provide [the service] to eligible patients. An "A" recommendation indicates good evidence that [the service] improves important health outcomes and that benefits substantially outweigh harms.

**Grade B** The panel recommends that clinicians routinely provide [the service] to eligible patients. A "B" recommendation indicates at least fair evidence that [the service] improves important health outcomes and concludes that benefits outweigh harms.

**Grade C** The panel recommends that clinicians routinely provide [the service] to eligible patients. A "C" recommendation indicates that there was consensus among the panel to recommend [the service] but that the evidence that [the service] is effective is lacking, of poor quality, or conflicting, or the balance of benefits and harms cannot be reliably determined from available evidence.

**Grade D** The panel recommends against clinicians routinely providing [the service]. A "D" recommendation indicates at least fair evidence that [the service] is ineffective or that harm outweighs benefit.

**Grade I** The panel concludes that the evidence is insufficient to recommend for or against [the service]. An "I" recommendation indicates that evidence that [the service] is effective is lacking, of poor quality, or conflicting, and the balance of

benefits and harms cannot be determined, and that the panel lacked a consensus to recommend it.

#### Net Benefit

The levels of net benefit are based on clinical assessment. Estimated net benefit may be downgraded based on uncertainty in estimates of benefits and harms.

Substantial Benefit: Benefit greatly outweighs harm

Moderate Benefit: Benefit outweighs harm

Small/weak Benefit: Benefit outweighs harm to a minimally clinically important degree

None/negative Benefit: Harms equal or outweigh benefit, less than clinically important

#### COST ANALYSIS

A formal cost analysis was not performed and published cost analyses were not reviewed.

#### METHOD OF GUIDELINE VALIDATION

Peer Review

#### DESCRIPTION OF METHOD OF GUIDELINE VALIDATION

After extensive review within the expert panel and executive committee, the guidelines were reviewed and approved by the American College of Chest Physicians (ACCP) Health and Science Policy Committee and then by the American College of Chest Physicians Board of Regents.

### RECOMMENDATIONS

#### MAJOR RECOMMENDATIONS

Each recommendation is rated based on the levels of evidence (good, fair, poor), net benefit (substantial, moderate, small/weak, none/negative), and the grades of the recommendations (A, B, C, D, I). Definitions are presented at the end of the "Major Recommendations" field.

1. All patients with known or suspected lung cancer should undergo a thorough history, physical examination, and standard laboratory tests as a screen for metastatic disease. Level of evidence, fair; benefit, substantial; grade of recommendation, B
2. Patients with lung cancer and a paraneoplastic syndrome should not be precluded from potentially curative therapy on the basis of these symptoms

alone. Level of evidence, fair; benefit, substantial; grade of recommendation, C

### Definitions:

#### Grades of Recommendations and Estimates of Net Benefit

The grade of the strength of recommendations is based on both the quality of the evidence and the net benefit of the service (i.e., test, procedure, etc).

**Grade A** The panel strongly recommends that clinicians routinely provide [the service] to eligible patients. An "A" recommendation indicates good evidence that [the service] improves important health outcomes and that benefits substantially outweigh harms.

**Grade B** The panel recommends that clinicians routinely provide [the service] to eligible patients. A "B" recommendation indicates at least fair evidence that [the service] improves important health outcomes and concludes that benefits outweigh harms.

**Grade C** The panel recommends that clinicians routinely provide [the service] to eligible patients. A "C" recommendation indicates that there was consensus among the panel to recommend [the service] but that the evidence that [the service] is effective is lacking, of poor quality, or conflicting, or the balance of benefits and harms cannot be reliably determined from available evidence.

**Grade D** The panel recommends against clinicians routinely providing [the service]. A "D" recommendation indicates at least fair evidence that [the service] is ineffective or that harm outweighs benefit.

**Grade I** The panel concludes that the evidence is insufficient to recommend for or against [the service]. An "I" recommendation indicates that evidence that [the service] is effective is lacking, of poor quality, or conflicting, and the balance of benefits and harms cannot be determined, and that the panel lacked a consensus to recommend it.

#### Net Benefit

The levels of net benefit are based on clinical assessment. Estimated net benefit may be downgraded based on uncertainty in estimates of benefits and harms.

**Substantial Benefit:** Benefit greatly outweighs harm

**Moderate Benefit:** Benefit outweighs harm

**Small/weak Benefit:** Benefit outweighs harm to a minimally clinically important degree

**None/negative Benefit:** Harms equal or outweigh benefit, less than clinically important.

#### CLINICAL ALGORITHM(S)

None provided

## EVIDENCE SUPPORTING THE RECOMMENDATIONS

### TYPE OF EVIDENCE SUPPORTING THE RECOMMENDATIONS

The type of supporting evidence is identified and graded for each recommendation (see "Major Recommendations").

## BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

### POTENTIAL BENEFITS

Asymptomatic patients and patients with symptoms related to the primary tumor have better 5-year survivals than those with systemic symptoms or symptoms indicating metastatic disease. Assessing symptoms, signs and standard laboratory tests in a standardized manner in patients presenting with known or suspected lung cancer may serve as a useful screen for identifying patients with a higher likelihood of metastatic disease.

### POTENTIAL HARMS

Not stated

## IMPLEMENTATION OF THE GUIDELINE

### DESCRIPTION OF IMPLEMENTATION STRATEGY

1. The American College of Chest Physicians (ACCP) is developing a set of PowerPoint slide presentations for physicians to download and use for physician and allied health practitioners education programs.
2. The ACCP is developing a Quick Reference Guide (QRG) in print and PDA formats for easy reference.

## INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

### IOM CARE NEED

Living with Illness

### IOM DOMAIN

Effectiveness

## IDENTIFYING INFORMATION AND AVAILABILITY

### BIBLIOGRAPHIC SOURCE(S)



Beckles MA, Spiro SG, Colice GL, Rudd RM. Initial evaluation of the patient with lung cancer: symptoms, signs, laboratory tests, and paraneoplastic syndromes. Chest 2003 Jan; 123(1 Suppl): 97S-104S. [54 references] [PubMed](#)

#### ADAPTATION

Not applicable: The guideline was not adapted from another source.

#### DATE RELEASED

2003 Jan

#### GUIDELINE DEVELOPER(S)

American College of Chest Physicians - Medical Specialty Society

#### GUIDELINE DEVELOPER COMMENT

The guideline development panel was composed of members and nonmembers of the American College of Chest Physicians (ACCP) who were known to have expertise in various areas of lung cancer management and care, representing multiple specialties from the following 13 national and international medical associations:

- Alliance for Lung Cancer Advocacy, Support, and Education (a patient support group)
- American Association for Bronchology
- American Cancer Society
- American College of Physicians
- American College of Surgeons Oncology Group
- American Society of Clinical Oncology
- American Society for Therapeutic Radiology and Oncology
- American Thoracic Society
- Association of Community Cancer Centers
- Canadian Thoracic Society
- National Comprehensive Cancer Network
- Oncology Nursing Society
- Society of Thoracic Surgeons

The specialties included pulmonary/respiratory medicine, critical care, medical oncology, thoracic surgery, radiation oncology, epidemiology, law, and medical ethics.

#### SOURCE(S) OF FUNDING

Funding for both the evidence reviews and guideline development was provided through an unrestricted educational grant from Bristol-Myers Squibb, which had no other role in the evidence review or guideline development process or content.

#### GUIDELINE COMMITTEE

## American College of Chest Physicians (ACCP) Expert Panel on Lung Cancer Guidelines

### COMPOSITION OF GROUP THAT AUTHORED THE GUIDELINE

Primary Authors: Michael A. Beckles, MB, BS; Stephen G. Spiro, MD; Gene L. Colice, MD, FCCP; Robin M. Rudd, MD

### FINANCIAL DISCLOSURES/CONFLICTS OF INTEREST

Information about potential conflicts of interest were collected from each member of the expert panel or writing committee at the time of their nomination in accordance with the policy of the American College of Chest Physicians. Information on conflicts of interest for each panelist is listed in the guideline.

### GUIDELINE STATUS

This is the current release of the guideline.

### GUIDELINE AVAILABILITY

Electronic copies: Available to subscribers of [Chest - The Cardiopulmonary and Critical Care Journal](#).

Print copies: Available from the American College of Chest Physicians, Products and Registration Division, 3300 Dundee Road, Northbrook IL 60062-2348.

### AVAILABILITY OF COMPANION DOCUMENTS

The following are available:

#### Background Articles:

- Alberts WM. Lung cancer guidelines. Introduction. Chest 2003 Jan;123(1 Suppl):1S-2S.
- McCrory DC, Colice GL, Lewis SZ, Alberts WM, Parker S. Overview of methodology for lung cancer evidence review and guideline development. Chest 2003 Jan;123(1 Suppl):3S-6S.
- Harpole LH, Kelley MJ, Schreiber G, Toloza EM, Kolimaga J, McCrory DC. Assessment of the scope and quality of clinical practice guidelines in lung cancer. Chest 2003 Jan;123(1 Suppl):7S-20S.
- Alberg AJ, Samet JM. Epidemiology of lung cancer. Chest 2003 Jan;123(1 Suppl):21S-49S.

Electronic copies: Available to subscribers of [Chest - The Cardiopulmonary and Critical Care Journal](#).

Print copies: Available from the American College of Chest Physicians, Products and Registration Division, 3300 Dundee Road, Northbrook IL 60062-2348.

## PATIENT RESOURCES

None available

## NGC STATUS

This NGC summary was completed by ECRI on June 30, 2003. The information was verified by the guideline developer on July 25, 2003.

## COPYRIGHT STATEMENT

This NGC summary is based on the original guideline, which is subject to the guideline developer's copyright restrictions.

© 1998-2004 National Guideline Clearinghouse

Date Modified: 11/8/2004

**FIRSTGOV**

